

Dispersive representation of $K \rightarrow 3\pi$ amplitudes and cusps*K. Kampf^{ab}, M. Knecht^c, J. Novotný^b, M. Zdráhal^{bd},^aPaul Scherrer Institut, CH-5232 Villigen PSI, Switzerland^bIPNP, MFF, Charles University, V Holešovičkách 2, CZ-180 00 Prague 8, Czech Republic^cCentre de Physique Théorique[†], CNRS-Luminy, Case 907, F-13288 Marseille Cedex 9, France^dFaculty of Physics, University of Vienna, Boltzmanngasse 5, A-1090 Vienna, Austria

The NA48/2 collaboration has shown clear experimental evidence for a cusp in the data for $K \rightarrow \pi\pi^0\pi^0$. This effect can be used to extract information on the $\pi\pi$ scattering lengths. We address this issue using a two-loop dispersive construction of $\pi\pi \rightarrow \pi\pi$ and $K \rightarrow \pi\pi\pi$ amplitudes in the presence of isospin breaking.

1. Introduction

The observation of a cusp anomaly in the $\pi^0\pi^0$ invariant mass distribution in the data on the $K^\pm \rightarrow \pi^\pm\pi^0\pi^0$ decay collected by the NA48/2 collaboration [1] has triggered some theoretical activity [2]-[5]. The basic explanation of the appearance of a unitarity cusp is very simple [2]: the amplitude for $K^+ \rightarrow \pi^+\pi^0\pi^0$ has two basic contributions, one of which corresponds to the $\pi^+\pi^-$ intermediate state rescattering to $\pi^0\pi^0$. This intermediate state in the s-channel generates (at the one-loop level) a square root singularity [6], and the corresponding amplitude behaves at this level for $s \sim 4m_+^2$ (m_+ is the mass of π^+) as

$$A(s) = R(\sigma_+^2) + \pi S(\sigma_+^2) \begin{cases} i\sigma_+, & s > 4m_+^2 \\ -\tilde{\sigma}_+, & s < 4m_+^2 \end{cases}, \quad (1)$$

where $\sigma_P = \sqrt{1 - 4m_P^2/s}$, $\tilde{\sigma}_P = \sqrt{4m_P^2/s - 1}$. The functions $R(\sigma_+^2)$ and $S(\sigma_+^2)$ can be expressed as convergent series in $s - 4m_+^2$ in the physical region of the $K^+ \rightarrow \pi^+\pi^0\pi^0$ decay. This singularity appears at $4m_+^2$, which is above the physi-

cal threshold $4m_0^2$ (m_0 is the mass of π^0), and the cusp is a result of the interference of the part containing the singularity and the rest without it. It is clear that the cusp appears only in the isospin breaking case and its strength is sensitive to the $\pi\pi$ scattering amplitude at the threshold. This is the reason why the investigation of the cusp effect can in principle serve as an independent method for the experimental determination of the $\pi\pi$ scattering lengths, provided a model independent description of the corresponding amplitude can be given.

In [3] Cabibbo and Isidori have proposed to use the assumed simple analytical properties of the amplitude ($\pi\pi$ amplitudes are considered in the form (1) with $R(\sigma_+^2)$ and $S(\sigma_+^2)$ polynomial) and the unitarity of the scattering matrix in order to express the amplitude near the threshold as an expansion in scattering lengths a_i . This idea was further investigated in [4]. Another possibility how to address this issue is the framework of non-relativistic effective field theory as developed in [5]. This is done as a combined expansion again in a_i , and also in the pion momenta. Both of these different approaches compute contributions to the K^+ decay amplitude up to order $O(a_i^2)$. Each of these approaches has its own limitations that result from the assumptions made. Moreover, they give no connection between cusp effects and the traditional PDG parameterization of the K decay amplitudes. We consider an alternative

*Presented by M. Z. and K. K. at Flavianet Kaon Workshop 08 and QCD 08. This work was supported in part by the Center for Particle Physics (project no. LC 527), GACR (grant no. 202/07/P249) and by the EU Contract No. MRTN-CT-2006-035482, "FLAVIANet".

In memoriam of our friend and colleague Jan Stern.

[†]Unité Mixte de Recherche (UMR 6207) du CNRS et des Universités Aix-Marseille 1 et 2 et Sud Toulon-Var, Laboratoire affilié à la FRUMAM (FR 2291)

approach, which rests on general properties, unitarity, analyticity, crossing symmetry, relativistic invariance, and chiral power counting for partial wave amplitudes. This leads, through a two-step iterative procedure, to a two-loop representation of the $K \rightarrow \pi\pi\pi$ amplitudes.

In the following we illustrate the method on the simplest case of the $K_L \rightarrow 3\pi^0$ decay, mainly because the analytical expressions are less involved, but also because, for the time being, we want to avoid addressing some further issues appearing in the treatment of the processes involving charged pions. The cusp in this decay has been observed by KTeV [7], as well as by NA48/2 [8].

2. Reconstruction theorem

The approach we wish to implement proceeds in parallel with the construction of the two-loop representation for the $\pi\pi$ scattering amplitude achieved in [9,10]. We shall use the fact that it can be extended to other processes and that isospin symmetry is not an essential ingredient [11]. Here we are interested in $K\pi \rightarrow \pi\pi$, related to $K \rightarrow 3\pi$ by crossing symmetry. The essential ingredients required in order to implement this construction are the following. First, we need a decomposition of the amplitude of the type

$$\mathcal{A}(s, t, u) = 16\pi(f_0(s) + 3f_1(s)\cos\theta) + \mathcal{A}_{\ell \geq 2} \quad (2)$$

with the following chiral behaviour,

$$\text{Re } \mathcal{A}_{\ell \geq 2} \sim O(p^4), \quad \text{Im } \mathcal{A}_{\ell \geq 2} \sim O(p^8), \quad (3)$$

$$\text{Re } f_\ell \sim O(p^2), \quad \text{Im } f_\ell \sim O(p^4), \quad \ell = 0, 1. \quad (4)$$

Having this, one can reconstruct the amplitude $\mathcal{A}(s, t, u)$ of a process $AB \rightarrow CD$ to $O(p^8)$ from the knowledge of the imaginary parts of S and P partial waves of all the crossed amplitudes:

$$\begin{aligned} \mathcal{A}(s, t, u) = & P(s, t; u) + \Phi_0(s) \\ & + [s(t - u) + (m_A^2 - m_B^2)(m_C^2 - m_D^2)] \Phi_1(s) \\ & + \text{crossed channels} + O(p^8), \end{aligned} \quad (5)$$

where $P(s, t; u)$ is a polynomial having the same s, t, u symmetries as the amplitude $\mathcal{A}(s, t; u)$ and of at most third order in the Mandelstam variables. Φ_0 and Φ_1 are the dispersive integrals of

the partial waves of the s -channel amplitude

$$\Phi_0(s) = 16s^3 \int_{\text{thresh.}}^{\infty} dx \frac{\text{Im } f_0(x)}{x^3(x - s)}, \quad (6)$$

$$\Phi_1(s) = 48s^3 \int_{\text{thresh.}}^{\infty} dx \frac{\text{Im } f_1(x)}{x^3(x - s) \lambda_{AB}^{1/2}(x) \lambda_{CD}^{1/2}(x)}$$

and similar for the t - and u -crossed channel [$\lambda_{AB}(x) = (s - (m_A + m_B)^2)(s - (m_A - m_B)^2)$].

The imaginary parts that enter the above expressions are obtained from the unitarity relation,

$$\begin{aligned} \text{Im } f_\ell^{i \rightarrow f}(s) = & \sum_k \frac{1}{S} \frac{\lambda_k^{1/2}(s)}{s} \\ & \times f_\ell^{i \rightarrow k}(s) \left(f_\ell^{f \rightarrow k}(s) \right)^* \theta(s - \text{thr}_k), \end{aligned} \quad (7)$$

projected on the corresponding partial waves. The sum goes over all the possible intermediate states k (S is a symmetry factor, $S = 2$ for indistinguishable states and $S = 1$ otherwise). In the low-energy region, and up to two-loops, these are restricted to pairs of light pseudoscalar mesons. To the extent that we are interested in the decay region only, we may further restrict them to intermediate $\pi\pi$ states. The contributions from other intermediate states, like *e.g.* $K\pi$, can be expanded in powers of the Mandelstam variables and absorbed into the polynomial $P(s, t, u)$.

This unitarity relation and the reconstruction theorem can be used iteratively, *i.e.* starting from the LO amplitudes, we obtain the NLO results. S and P partial wave projections thereof then allow to obtain the NNLO expressions. Details of the first step will be presented in the next section.

According to the reconstruction theorem and due to the crossing symmetry, the two-loop representation of the $K_L \rightarrow 3\pi^0$ amplitude looks like:

$$\begin{aligned} \mathcal{A}_{L;00}(s, t, u) = & P_{L;00} + \Phi_0^{L;00}(s) + \Phi_0^{L;00}(t) \\ & + \Phi_0^{L;00}(u) + O(p^8) \end{aligned} \quad (8)$$

with the polynomial

$$\begin{aligned} P_{L;00} = & C_F (A_{00}^L M_K^2 + \{C_{00}^L [(s - s_0^L)^2] \\ & + E_{00}^L [(s - s_0^L)^3]\} + \{s \leftrightarrow t\} + \{s \leftrightarrow u\}), \end{aligned} \quad (9)$$

where the centre of Dalitz plot was defined as $s_0^L = 1/3 M_K^2 + m_0^2$ and C_F corresponds to the standard normalization, $C_F = -\frac{3}{5} V_{us}^* V_{ud} \frac{G_F}{\sqrt{2}}$.

3. First iteration: one-loop expressions

As already stated, we need the leading order $\pi\pi$ and $K\pi \rightarrow \pi\pi$ scattering amplitudes. From the chiral perturbation theory we know that at $O(p^2)$ they are represented by polynomials of at most first order in the Mandelstam variables. Their particular choice (connected also with the particular choice of the polynomial of the reconstruction theorem) is important since different choices can possibly lead to different convergence properties of the chiral expansion and affect the stability of the fit to the data. The standard choice (believed to be stable) is the expansion in sub-threshold parameters. For the $\pi\pi$ amplitude, this corresponds to

$$\begin{aligned}\mathcal{A}_{\text{LO}}^{+-;00} &= -\frac{\beta_{\pm 0}}{F_\pi^2} \left(s - \frac{2}{3}m_+^2 - \frac{2}{3}m_0^2 \right) - \frac{\alpha_{\pm 0}m_0^2}{3F_\pi^2}, \\ \mathcal{A}_{\text{LO}}^{00;00} &= \frac{\alpha_{00}m_0^2}{F_\pi^2}.\end{aligned}\quad (10)$$

Another possibility would consist in choosing the scattering length and effective range parameter as independent coefficient. It is even possible to adjust the polynomial part $P(s, t; u)$ of the $\pi\pi$ amplitude so that these coefficients retain their physical interpretation up to two loops, just as in the non-relativistic approach [5]. We shall study this option elsewhere [12]. In the case of $K_L\pi^0 \rightarrow \pi^0\pi^0$ and $K_L\pi^0 \rightarrow \pi^+\pi^-$ amplitudes we have

$$\begin{aligned}\mathcal{A}_{L;00}^{\text{LO}} &= C_F A_{00}^L M_K^2, \\ \mathcal{A}_{L;+-}^{\text{LO}} &= C_F [B_{+-}^L (s - s_\pm^L) + A_{+-}^L M_K^2],\end{aligned}\quad (11)$$

where $s_\pm^L = (M_K^2 + m_0^2 + 2m_\pm^2)/3$.

The $O(p^2)$ chiral perturbation theory result is reproduced by special values of the parameters ($\alpha_{00} = 1$, $\beta_{\pm 0} = 1$, $\alpha_{\pm 0} = (2m_+^2 - m_0^2)/m_0^2$; cf. [13]) and similarly for the $K\pi$ part (see [14]).

Using these amplitudes in the first iteration, we obtain the one-loop result for $\Phi_0^{L;00}$ of (8) as

$$\begin{aligned}\Phi_0^{L;00}(s) &= \frac{C_F}{2F_\pi^2} A_{00}^L M_K^2 \alpha_{00} m_0^2 \bar{J}_0(s) \\ &\quad - \frac{C_F}{F_\pi^2} [\beta_{\pm 0} (s - \frac{2}{3}m_+^2 - \frac{2}{3}m_0^2) + \frac{1}{3}\alpha_{\pm 0}m_0^2] \\ &\quad \times [A_{+-}^L M_K^2 + B_{+-}^L (s - s_{+-}^L)] \bar{J}_\pm(s) \\ &\quad + \text{polynomial} + O(p^6).\end{aligned}\quad (12)$$




Figure 1. The partial decay rate $K_L \rightarrow 3\pi^0$ (in arbitrary units) as a function of the invariant mass of the $\pi^0\pi^0$ pair squared for one particular choice of parameters. Within the small frame the cusp region is zoomed.

The loop functions $\bar{J}$ are defined by

$$\bar{J}_{PQ}(s) = \frac{s}{16\pi^2} \int_{(m_P+m_Q)^2}^{\infty} dx \frac{\lambda_{PQ}^{1/2}(x)}{x(x-s)}, \quad (13)$$

which simplifies for both masses the same to

$$\bar{J}_P(s) = \frac{1}{16\pi^2} \left(2 + \sigma_P \ln \frac{\sigma_P - 1}{\sigma_P + 1} \right). \quad (14)$$

Our NLO parameterization of the amplitude (8) with $\Phi_0^{L;00}$ given by (12) can be now fitted to experimental data, and so we can get the $O(p^2)$ values of all the seven constants appearing in the LO parameterization of (10) and (11). It is also important to have in mind that our parameterization of the amplitude encompasses the chiral perturbation theory result [14] as a particular case for special values of the parameters.

As an illustration of the cusp effect in the $K_L \rightarrow 3\pi^0$ decay, we plot the resulting partial decay rate for one particular choice of the sub-threshold parameters in the Fig. 1.

4. Second iteration: some remarks

The second iteration leads to the two-loop expression of the $K \rightarrow 3\pi$ amplitude. However, a

few complications have to be dealt with. As we have seen above, in order to get the discontinuities of the function $\Phi_0^{L;00}(x)$, we need an appropriate analytic continuation of the unitarity relation (and therefore also an analytic continuation of the S and P partial wave projections of the amplitudes obtained by the first iteration) below the physical threshold. In the isospin limit the solution is well known. The general method developed in [15] is based on the careful definition of the t -integration contour $C(t_+, t_-)$ in the formula for the partial wave projections, where the endpoints are

$$t_{\pm}(s) = \frac{1}{2} \left(3s_0^L - s \pm \lambda_{L0}^{1/2}(s)\sigma_0 \right) + i\varepsilon, \quad (15)$$

$$\text{sign } \varepsilon = \text{sign } \frac{\partial t_{\pm}(s)}{\partial M_K^2}. \quad (16)$$

The contour $C(t_+, t_-)$ is defined in such a way that it avoids the intersection with the cuts attached to branch points corresponding to the normal threshold of the amplitude $\mathcal{A}_{L;00}(s, t, u)$ in the t - and u - channel. The trajectory of $t_{\pm}(s)$ and the basic types of the contour $C(t_+, t_-)$ in the complex t -plane are depicted in [15]. In the case of the $K_L\pi^0 \rightarrow \pi^0\pi^0$ and $K_L\pi^0 \rightarrow \pi^+\pi^-$ scattering, the generalization of this prescription beyond the isospin limit is straightforward. However, this is not the case of *e.g.* the experimentally more interesting process $K^+\pi^- \rightarrow \pi^0\pi^0$, where the naive application of the prescription [15] shows some problems. Namely, for the reconstruction of the amplitude of this process beyond NLO we need to compute the partial waves of $K^+\pi^- \rightarrow \pi^+\pi^-$, where the trajectory of $t_-(s)$ crosses the t -channel cut of this amplitude instead of avoiding it, contrary to the isospin limit. In addition, the two-loop amplitude for $K^+\pi^- \rightarrow \pi^0\pi^0$ suffers from (complex) anomalous threshold stemming from the triangle Landau singularity. The consequence is that a careful analytic continuation of the normal dispersive integrals entering the representation of the amplitude by means of the reconstruction theorem has to be performed. These technical issues deserve a detailed further discussion which is however beyond the scope of this talk.

5. Conclusion

The cusp effect in $K \rightarrow 3\pi$ offers an interesting possibility to extract quantitative information on $\pi\pi$ scattering lengths from the experimental data. We have outlined a construction, in a fully relativistic framework, of the corresponding two-loop amplitudes, based on general properties, unitarity, analyticity, crossing, and chiral counting for the partial waves. Our analysis provides two parameterizations, one in terms of the subthreshold parameters and the other, as in existing analyses, directly in terms of scattering lengths.

REFERENCES

1. J. R. Batley *et al.* [NA48/2 Collaboration], Phys. Lett. B **633** (2006) 173.
2. N. Cabibbo, Phys. Rev. Lett. **93** (2004) 121801.
3. N. Cabibbo and G. Isidori, JHEP **0503** (2005) 021.
4. E. Gamiz, J. Prades and I. Scimemi, Eur. Phys. J. C **50** (2007) 405.
5. G. Colangelo, J. Gasser, B. Kubis and A. Rusetsky, Phys. Lett. B **638** (2006) 187; M. Bissegger *et al.*, Phys. Lett. B **659** (2008) 576 and arXiv:0807.0515 [hep-ph].
6. U. G. Meissner, G. Muller and S. Steininger, Phys. Lett. B **406** (1997) 154.
7. E. Abouzaid *et al.* [KTeV Collaboration], Phys. Rev. D **78** (2008) 032009.
8. See *e.g.* G. Lamanna at QCD08.
9. J. Stern, H. Sazdjian and N. H. Fuchs, Phys. Rev. D **47** (1993) 3814.
10. M. Knecht, B. Moussallam, J. Stern and N. H. Fuchs, Nucl. Phys. B **457** (1995) 513.
11. M. Zdráhal and J. Novotný, arXiv:0806.4529 [hep-ph].
12. K. Kampf, M. Knecht, J. Novotný, M. Zdráhal: *in preparation*.
13. M. Knecht and R. Urech, Nucl. Phys. B **519** (1998) 329.
14. J. Bijmans and F. Borg, Nucl. Phys. B **697** (2004) 319.
15. J. B. Bronzan and C. Kacser, Phys. Rev. **132** (1963) 2703; V. V. Anisovich and A. A. Anselm, Usp. Fyz. Nauk **88** (1966) 287.